

bsm-55571R**[Primary Antibody]**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

FSTL3 Recombinant Rabbit mAb**— DATASHEET —**


Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:50-1:200) IHC-F (1:50-1:200) IF (1:50-1:200) ICC/IF (1:50-1:200) Reactivity: Human Predicted MW.: 28 kDa Subcellular Location: Cytoplasm ,Nucleus
Clonality: Recombinant	CloneNo.: 9F5	
GeneID: 10272	SWISS: O95633	
Target: FSTL3		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.22μm filtered solution in 20mM hepes7.4, 150mm NaCl, 50% glycerin,0.03% proclin. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: FLRG (follistatin-related gene, follistatin-like-3 or FSTL3) is a member of the follistatin-module protein family, which is composed of extracellular matrix-associated glycoproteins thought to act in a paracrine manner to bind morphogens or growth/differen-tiation factors and regulate their activity during development. The FSTL3 protein contains 2 potential N-glycosylation sites and the predicted mass of the unmodified core protein is 27 kDa. FLRG is expressed in a wide range of human and murine adult tissues and its expression seems to be tightly regulated during murine embryogenesis. Immunohistochemistry reveals the presence of FLRG in the basement membrane between the dermis and the epidermis and around blood vessels.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded Human Heart; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with FSTL3 Monoclonal Antibody, Unconjugated(bsm-55571R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Placenta; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with FSTL3 Monoclonal Antibody, Unconjugated(bsm-55571R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



4% Paraformaldehyde-fixed HeLa (H) cell; Triton X-100 at r.t. for 20 min; Antibody incubation with (FSTL3) monoclonal Antibody, unconjugated (bsm-55571R) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-40295G-FITC) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.