bs-19079R

[Primary Antibody]


NDUFB1 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –––––		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500) IF (1:100-500)
GenelD: 4707	SWISS: 075438	(,)
Target: NDUFB1		Reactivity: Human
Immunogen: KLH conjugated syr 21-58/58.	nthetic peptide derived from human NDUF	FB1:
Purification: affinity purified by Protein A		Predicted 7 kDa
Concentration: 1mg/ml		MW.: *****
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Cytoplasm Location:
localizes to the mat member of the com encoded by a gene Chromosome 14 ho of the human geno (PSEN1) gene, whic the development o is also located on c the genetic disorde	ino acid single-pass membrane protein the trix side of the mitochondrial membrane. A pplex I NDUFB1 subunit family, NDUFB1 is that maps to human chromosome 14q32. buses over 700 genes and comprises nearly me. Chromosome 14 encodes the presinil is one of the three key genes associated f Alzheimer's disease (AD). The SERPINA1 thromosome 14 and, when defective, leads or ?-antitrypsin deficiency, which is evere lung complications and liver dysfunc	A 12. y 3.5% in 1 with gene s to
- VALIDATION IMAGES		

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NDUFB1) Polyclonal Antibody, Unconjugated (bs-19079R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.