## bs-5935R

## [ Primary Antibody ]

# Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# **KLHL7 Rabbit pAb**

- DATASHEET -

**Host:** Rabbit **Isotype:** IgG

Clonality: Polyclonal

**GenelD:** 55975 **SWISS:** Q8IXQ5

Target: KLHL7

**Immunogen:** KLH conjugated synthetic peptide derived from human KLHL7:

51-150/856.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** This gene encodes a BTB-Kelch-related protein. The encoded

protein may be involved in protein degradation. Mutations in this gene have been associated with retinitis pigmentosa 42. [provided

by RefSeq, Feb 2010].

**Applications: IHC-P** (1:100-500)

**IHC-F** (1:100-500) **IF** (1:100-500)


Reactivity: Mouse, Rat

(predicted: Human, Pig, Cow, Chicken, Dog, Horse)


Predicted MW.: 64 kDa

Subcellular Location: Nucleus

### VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KLHL7) Polyclonal Antibody, Unconjugated (bs-5935R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KLHL7) Polyclonal Antibody, Unconjugated (bs-5935R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.