

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## **Recombinant Mouse HSD11B2 Protein, N-His**

Catalog Number:	bs-105563P
Species:	Mouse
AA Seq:	82-331/386
Predicted MW:	29.35 kDa
Tags:	N-His
Activity:	Not tested
Purity:	>90% as determined by SDS-PAGE.
Purification:	AC
Form:	Lyophilized
Storage:	Lyophilized from a solution in PBS pH 7.4, 0.02% NLS, 1mM EDTA, 4% Trehalose, 1%
	Mannitol.
	Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 2 to 8°C for
	frequent use. Store at -20 to -80°C for twelve months from the date of receipt.
Background:	There are at least two isozymes of the corticosteroid 11-beta-dehydrogenase, a microsomal
	enzyme complex responsible for the interconversion of cortisol and cortisone. The type I
	isozyme has both 11-beta-dehydrogenase (cortisol to cortisone) and 11-oxoreductase
	(cortisone to cortisol) activities. The type II isozyme, encoded by this gene, has only 11-beta-
	dehydrogenase activity. In aldosterone-selective epithelial tissues such as the kidney, the
	type II isozyme catalyzes the glucocorticoid cortisol to the inactive metabolite cortisone,
	thus preventing illicit activation of the mineralocorticoid receptor. In tissues that do not
	express the mineralocorticoid receptor, such as the placenta and testis, it protects cells from
	the growth-inhibiting and/or pro-apoptotic effects of cortisol, particularly during embryonic
	development. Mutations in this gene cause the syndrome of apparent mineralocorticoid
	excess and hypertension. [provided by RefSeq, Feb 2010]