

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Recombinant Human CD25/IL2RA Protein, C-His

| Catalog Number: | bs-105714P |
|-----------------|---|
| Species: | Human |
| AA Seq: | 1-240/272 |
| Predicted MW: | 28.16 kDa |
| Tags: | C-His |
| Purity: | >90% as determined by SDS-PAGE. |
| Purification: | AC |
| Form: | Lyophilized |
| Storage: | Lyophilized from a solution in PBS pH 7.4, 1mM EDTA, 4% Trehalose, 1% Mannitol. |
| | Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 2 to 8°C for |
| | frequent use. Store at -20 to -80°C for twelve months from the date of receipt. |
| Background: | The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains, together with the |
| | common gamma chain (IL2RG), constitute the high-affinity IL2 receptor. Homodimeric alpha |
| | chains (IL2RA) result in low-affinity receptor, while homodimeric beta (IL2RB) chains |
| | produce a medium-affinity receptor. Normally an integral-membrane protein, soluble IL2RA |
| | has been isolated and determined to result from extracellular proteolyisis. Alternately- |
| | spliced IL2RA mRNAs have been isolated, but the significance of each is presently unknown. |
| | Mutations in this gene are associated with interleukin 2 receptor alpha deficiency. Patients |
| | with severe Coronavirus Disease 2019 (COVID-19), the disease caused by the novel severe |
| | acute respiratory syndrome coronavirus 2 (SARS-CoV-2), have significantly elevated levels of |
| | IL2R in their plasma. Similarly, serum IL-2R levels are found to be elevated in patients with |
| | different types of carcinomas. Certain IL2RA and IL2RB gene polymorphisms have been |
| | associated with lung cancer risk. [provided by RefSeq, Jul 2020] |
| | |