

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## FAK Recombinant Rabbit mAb

Catalog Number: bsm-52277R

Target Protein: FAK

Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Recombinant

Clone No.: 2F4

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), Flow-Cyt (1:50-100),

ICC/IF (1:50-200)

Reactivity: Human, Mouse, Rat (predicted:Human, Mouse, Rat)

Predicted MW: 116 kDa

Subcellular Cell membrane, Cytoplasm, Nucleus

Locations:

Entrez Gene: 5747 Swiss Prot: Q05397

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Non-receptor protein-tyrosine kinase implicated in signaling pathways involved in cell

motility, proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody cross-linking, or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid, or via LDL receptor occupancy. Plays a potential role in oncogenic transformations resulting in increased kinase activity. [SUBCELLULAR LOCATION] Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Note=Constituent of focal

adhesions.