

Recombinant human CD80 protein, C-His (HEK293)

Catalog Number: bs-43750P

Concentration: >0.5mg/ml

Species: Human

AA Seq: 35-242/288

Predicted MW: 25.8

Tags: C-His

Endotoxin: Not analyzed

Purity: >90% as determined by SDS-PAGE

Purification: AC

Form: Liquid

Storage: PBS (pH=7.4)

Stored at -70°C or -20°C. Avoid repeated freeze/thaw cycles.

Background: CD80 is a member of the Ig superfamily and serves as the ligand for two T cell molecules, CD28 and CTLA4. Interactions between CD28 and CD80 on activated B cells result in enhanced T cell activation. CD80 is rapidly induced on the surface of in vitro activated B cells, Epstein Barr Virus (EBV) transformed B cell lines, Burkitts lymphoma cell lines, freshly isolated follicular B lymphoma cells, T cells, and monocytes. It is also expressed at high levels in dendritic cells. It reacts weakly with a small proportion of non activated normal B cells and with HTLV1 infected T cells. CD80 does not react with peripheral monocytes, resting and activated normal T cells, T cell lines and T cell clones, nor with myelomonocytic cell lines.

VALIDATION IMAGES



The purity of the protein is greater than 90% as determined by reducing SDS-PAGE.