
Recombinant human FADD Protein, N-His

Catalog Number: bs-42346P

Concentration: >1mg/ml

Species: Human

AA Seq: 1-208/208

Predicted MW: 26.9 kDa

Tags: N-His

Endotoxin: Not analyzed

Purity: >90% as determined by SDS-PAGE

Purification: AC


Form: Liquid

Storage: 20mM Tris-HCl (pH=8.0) with 4M Urea and 150mM NaCl

Stored at -70°C or -20°C. Avoid repeated freeze/thaw cycles.

Background: Predicted to enable several functions, including caspase binding activity; death effector domain binding activity; and tumor necrosis factor receptor superfamily binding activity. Involved in several processes, including hematopoietic or lymphoid organ development; negative regulation of activation-induced cell death of T cells; and positive regulation of CD8-positive, alpha-beta cytotoxic T cell extravasation. Acts upstream of or within extrinsic apoptotic signaling pathway in absence of ligand; motor neuron apoptotic process; and regulation of programmed cell death. Predicted to be located in several cellular components, including cell body; cytosol; and membrane raft. Predicted to be part of CD95 death-inducing signaling complex and ripoptosome. Predicted to be active in cytoplasm. Is expressed in several structures, including alimentary system; brain; genitourinary system; hemolymphoid system gland; and liver and biliary system. Human ortholog(s) of this gene implicated in leukemia. Orthologous to human FADD (Fas associated via death domain). [provided by Alliance of Genome Resources, Apr 2022]

VALIDATION IMAGES



The purity of the protein is greater than 90% as determined by reducing SDS-PAGE.