

**bs-0857R****[ Primary Antibody ]****beta-arrestin 1 + 2 Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 408**SWISS:** P49407**Target:** beta-arrestin 1 + 2**Immunogen:** KLH conjugated synthetic peptide derived from human Beta-arrestin 1: 241-340/418.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** Beta Arrestin 1 is a member of a family of proteins that are widely expressed but especially abundant in the central nervous system. Serving as an adaptor or scaffold molecule, beta Arrestin 1 is essential for mitogenic signaling. It mediates agonist dependent desensitization and internalization of G protein coupled receptors (GPCRs, e.g., beta 2 adrenergic receptor). After binding to their ligand and interacting with heterotrimeric G proteins, GPCRs are phosphorylated by G protein receptor kinases (GRKs) on serine residues. Beta Arrestin 1 has important roles in the cytoplasm and at the plasma membrane in the desensitization and internalization of G protein coupled receptors (GPCRs) and is increasingly appreciated to play an important role in the endocytosis and signaling of GPCRs. Beta Arrestin 1 in the cytosol is phosphorylated by ERK1 and 2 on serine 412 in a negative feedback mechanism and binds to the phosphorylated receptors at the plasma membrane. Serine 412 is then dephosphorylated and the GPCRs are internalized, leading to activation of the Ras, Raf, ERK1 and 2 signaling pathway.

**Applications:** WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)

**Reactivity:** Human, Mouse  
(predicted: Rat, Rabbit, Pig,  
Cow, Chicken, Dog,  
GuineaPig, Horse)

**Predicted**  
**MW.:** 45 kDa


**Subcellular** Cell membrane ,Cytoplasm  
**Location:** ,Nucleus

**— VALIDATION IMAGES —**

Sample: Lung (Mouse) Lysate at 40 ug Primary:  
Anti- beta-arrestin 1 + 2 (bs-0857R) at 1/1000  
dilution Secondary: IRDye800CW Goat Anti-  
Rabbit IgG at 1/20000 dilution Predicted band  
size: 45 kD Observed band size: 52 kD



Sample:Lung (Mouse) Lysate at 30 ug Primary:  
Anti-Beta-arrestin 1 (bs-0857R) at 1/300 dilution  
Secondary: IRDye800CW Goat Anti-Mouse IgG at  
1/20000 dilution Predicted band size: 45 kD  
Observed band size: 45 kD



Paraformaldehyde-fixed, paraffin embedded  
(human colon carcinoma); Antigen retrieval by  
boiling in sodium citrate buffer (pH6.0) for  
15min; Block endogenous peroxidase by 3%  
hydrogen peroxide for 20 minutes; Blocking  
buffer (normal goat serum) at 37°C for 30min;  
Antibody incubation with (beta-arrestin 1 + 2)  
Polyclonal Antibody, Unconjugated (bs-0857R)  
at 1:200 overnight at 4°C, followed by operating  
according to SP Kit(Rabbit) (sp-0023)  
instructions and DAB staining.