
Recombinant human Tissue Factor protein (Active, CHO)

Catalog Number: bs-48088P

Species: Human

AA Seq: 33-251/295

Predicted MW: 24.8

Tags: Tag free

Activity: Yes

Endotoxin: ≤ 10 EU/mg

Purity: $\geq 95\%$ as determined by SDS-PAGE.

Purification: AC

Form: Lyophilized

Storage: Lyophilized from a 0.22 μ m filtered solution containing PBS, 5% Mannitol and 0.01% Tween 80, pH7.4

Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 36 months at -20°C to -80°C in lyophilized state. 6 months at -20°C to -80°C under sterile conditions after reconstitution. 7-10 days at 2°C to 8°C under sterile conditions after reconstitution.

Background: This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.[provided by RefSeq, May 2010]