

bs-15179R**[Primary Antibody]****C3orf58 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn



sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 205428</p> <p>Target: C3orf58</p> <p>Immunogen: KLH conjugated synthetic peptide derived from human C3orf58: 351-430/430.</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: C3orf58 (chromosome 3 open reading frame 58), also known as DIA1, is a 430 amino acid secreted protein that belongs to the UPF0672 family. C3orf58 is encoded by a gene that maps to human chromosome 3q24. Chromosome 3 is made up of approximately 214 million bases encoding over 1,100 genes. Notably, there is a chemokine receptor gene cluster and a variety of human cancer related loci on chromosome 3. Particular regions of the chromosome 3 short arm are deleted in many types of cancer cells. Key tumor suppressing genes on chromosome 3 encode apoptosis mediator RASSF1, cell migration regulator HYAL1 and angiogenesis suppressor SEMA3B. Marfan Syndrome, porphyria, von Hippel-Lindau syndrome, osteogenesis imperfecta and Charcot-Marie-Tooth disease are a few of the numerous genetic diseases associated with chromosome 3.</p>	<p>Isotype: IgG</p> <p>SWISS: Q8NDZ4</p>	<p>Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)</p> <p>Reactivity: Human, Mouse, Rat (predicted: Cow, Chicken)</p> <p>Predicted MW.: 49 kDa</p> <p>Subcellular Location: Secreted ,Cytoplasm</p>
--	--	--

— VALIDATION IMAGES —

Sample: Lane 1: Mouse Colon tissue lysates Lane 2: Mouse Cerebrum tissue lysates Lane 3: Rat Colon tissue lysates Lane 4: Human 293T cell lysates Lane 5: Human U251 cell lysates Primary: Anti-C3orf58 (bs-15179R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 49 kDa Observed band size: 45 kDa

Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C3orf58) Polyclonal Antibody, Unconjugated (bs-15179R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.