

**bs-8485R**

**[ Primary Antibody ]**

## ZNF180 Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

### — DATASHEET —

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>ELISA</b> (1:5000-10000)
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 7733	<b>SWISS:</b> Q9UJW8	
<b>Target:</b> ZNF180		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human ZNF180: 501-600/692.		
<b>Purification:</b> affinity purified by Protein A		<b>Reactivity:</b> (predicted: Human, Rat)
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		<b>Predicted MW.:</b> 79 kDa
<b>Background:</b> Zinc finger proteins have been shown to interact with nucleic acids and to have diverse functions. The zinc finger domain is a conserved amino acid sequence motif containing 2 specifically positioned cysteines and 2 histidines that are involved in coordinating zinc. Kruppel-related proteins form 1 family of zinc finger proteins. See MIM 604749 for additional information on zinc finger proteins.[supplied by OMIM, Jul 2002]		<b>Subcellular Location:</b> Nucleus

### — SELECTED CITATIONS —

- **[IF=3.8]** Liang Xu. et al. Zinc finger protein 180 induces an apoptotic phenotype by activating METTL14 transcriptional activity in colorectal cancer. ONCOL REP. 2024 Oct;52(4):1-18 WB ;Human. 39054954