## bs-8485R

## [ Primary Antibody ]

## ZNF180 Rabbit pAb



sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	<b>Isotype:</b> IgG	Applications: WB (1:500-2000) ELISA (1:5000-10000)
Clonality: Polyclonal		
<b>GeneID:</b> 7733	SWISS: Q9UJW8	<b>Reactivity:</b> (predicted: Human, Rat)
Target: ZNF180		
Immunogen: KLH conjugated syn 501-600/692.	nthetic peptide derived from human ZNF	
Purification: affinity purified by Protein A		Predicted <sub>79 kDa</sub> MW.:
Concentration: 1mg/ml		Subcellular
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Nucleus
and to have diverse conserved amino a positioned cysteine coordinating zinc. I finger proteins. See	have been shown to interact with nucle functions. The zinc finger domain is a cid sequence motif containing 2 specific s and 2 histidines that are involved in Kruppel-related proteins form 1 family o MIM 604749 for additional information oplied by OMIM, Jul 2002]	ally f zinc

## 

• [IF=3.8] Liang Xu. et al. Zinc finger protein 180 induces an apoptotic phenotype by activating METTL14 transcriptional

activity in colorectal cancer. ONCOL REP. 2024 Oct;52(4):1-18 WB ;Human. 39054954