

bs-16194R

[Primary Antibody]

Fumarylacetoacetate hydrolase Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

| | | |
|---|----------------------|--|
| Host: Rabbit | Isotype: IgG | Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Pig) Predicted MW.: 47 kDa Subcellular Location: Secreted ,Extracellular matrix ,Cytoplasm |
| Clonality: Polyclonal | | |
| GeneID: 2184 | SWISS: P16930 | |
| Target: Fumarylacetoacetate hydrolase | | |
| Immunogen: KLH conjugated synthetic peptide derived from human Fumarylacetoacetate hydrolase: 21-120/419. | | |
| Purification: affinity purified by Protein A | | |
| Concentration: 1mg/ml | | |
| Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. | | |
| Background: This gene encodes the last enzyme in the tyrosine catabolism pathway. FAH deficiency is associated with Type 1 hereditary tyrosinemia (HT). [provided by RefSeq, Jul 2008] | | |

— SELECTED CITATIONS —

- **[IF=2.634]** Gao M et al. Efficient Generation of an Fah/Rag2 Dual-Gene Knockout Porcine Cell Line Using CRISPR/Cas9 and Adenovirus. DNA Cell Biol. 2019 Feb 14. ICC,IF ;Porcine. 30762444