

bs-18051R**[Primary Antibody]****HMG2L1 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Mouse, Rat (predicted: Human, Sheep, Cow, Dog, Horse) Predicted MW.: 66 kDa Subcellular Location: Nucleus
Clonality: Polyclonal		
GeneID: 10042	SWISS: Q9UGU5	
Target: HMG2L1		
Immunogen: KLH conjugated synthetic peptide derived from human HMG2L1: 1-100/601.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: High mobility group (HMG) proteins are nonhistone chromosomal proteins. See HMG2 (MIM 163906) for additional information on HMG proteins.[supplied by OMIM, Nov 2010]		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HMG2L1) Polyclonal Antibody, Unconjugated (bs-18051R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat lung); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HMG2L1) Polyclonal Antibody, Unconjugated (bs-18051R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.