- DATASHEET -

[Primary Antibody]

GlyR alpha 1 + 2 + 3 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DITITIONEL	1			
Host:	Rabbit	Isotype: IgG	Applications: WB (1:500-2000))
Clonality:	Polyclonal		IHC-P (1:100-50)0))0)
GeneID:	2741	SWISS: P23415	IF (1:100-500)	,0,
Target:	GlyR alpha 1 + 2 + 3	Reactivity: Human, Mouse, R	. R	
Immunogen:	mmunogen: KLH conjugated synthetic peptide derived from human GlyR alpha 1: 101-200/449. (predicted: Ra Cow, Chicken			
Purification:	affinity purified by Protein A			
Concentration:	1mg/ml		Predicted MW.: ^{49 kDa}	
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.			Subcellular Location: Cell membrane	ļ
Background:	Glycine receptors are memb superfamily, which mediate receptors are pentameric m	pers of the ligand-gated ion channel fast inhibitory neurotransmission. The embrane proteins which form chloride		

channels. Binding of glycine to its receptor produces an increase in chloride conductance and membrane hyperpolarisation. Four genes encoding glycine receptor alpha subunits have been identified, together with a single beta polypeptide. Each subunit consists of a large extracellular N-terminal region, four transmembrane domains, and a large cytoplasmic domain.

- VALIDATION IMAGES -



Sample: Lane 1: Mouse Cerebrum tissue lysates Lane 2: Mouse Cerebellum tissue lysates Lane 3: Mouse Testis tissue lysates Lane 4: Mouse Lung tissue lysates Lane 5: Rat Cerebrum tissue lysates Lane 6: Rat Cerebellum tissue lysates Lane 7: Rat Testis tissue lysates Lane 8: Human HeLa cell lysates Lane 9: Human A431 cell lysates Lane 10: Human SH-SY5Y cell lysates Primary: Anti-GlyR alpha 1 + 2 + 3 (bs-1524R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 49 kDa Observed band size: 53 kDa

	Sinalond ord
100 — 75 —	
63	
48 —	
35 —	
25 —	

Sample: Spinal cord (Mouse) Lysate at 40 ug Spinal cord (Rat) Lysate at 40 ug Primary: Anti-GlyR alpha 1 + 2 + 3 (bs-1524R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 49 kD Observed band size: 54 kD

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-GlyR alpha 1/2/3 Polyclonal Antibody, Unconjugated(bs-1524R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

0-500) Mouse, Rat ted: Rabbit, Pig, nicken, Dog, Horse)