bs-0358D-RBITC

[Secondary Antibodies]

Donkey Anti-Guinea Pig IgG H&L, RBITC conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

| - DATASHEET | | 400-901-9800 |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|
| Host: Donkey Clonality: Polyclonal | Isotype: IgG | Applications: IF (1:100-1000) Flow-Cyt (1:100-1000) |
| Target: Donkey Anti-Guinea Pig IgG H&L Purification: affinity purified by Protein G | | ICC/IF (1:100-1000) Excitation Spectrum: 540nm Emission spectrum: 574nm Reactivity: Guinea Pig |
| glycerol. | with 1% BSA, 0.03% Proclin300 and 50 | 0% |
| serum with normal l is important for our molecules are produ adaptive immune re functions; to bind to recruit other cells ar variability of the IgG | IgG), is one of the most abundant prot evels between 8-17 mg/mL in adult blo defence against microorganisms and t iced by B lymphocytes as a part of our sponse. The IgG molecule has two sep the pathogen that elicited the respon id molecules to destroy the antigen. Th pool is generated by somatic recombi pecificities in an individual at a given t b be 1011 variants. | bod. IgG the varate se and to he ination |