

bs-8337R**[Primary Antibody]****PANK1 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 53354	SWISS: Q8TE04	IHC-F (1:100-500)
Target: PANK1		IF (1:50-200)
Immunogen: KLH conjugated synthetic peptide derived from human PANK1: 501-598/598.		Reactivity: Mouse, Rat (predicted: Human, Pig, Sheep, Cow, Dog)
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Predicted MW.: 64 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cytoplasm
Background: The pantothenate kinase (PANK) family of proteins catalyzes the first step in coenzyme A (CoA). Pantothenate kinase 1 (PANK1) is a 598 amino acid member of the pantothenate kinase family that plays a role in the physiological regulation of the intracellular CoA concentration. Localized to the cytoplasm, PANK1 is strongly inhibited by acetyl-CoA and manyl-CoA, as well as by high concentration of non-esterified CoA (CoASH). Four known isoforms of PANK1 exist as a result of alternative splicing events. Of these isoforms, PANK1a and PANK1b have been identified as the catalytically active isoforms. Isoform PANK1a is most highly expressed in brain, heart, kidney, liver, skeletal muscle and kidney. Isoform PANK1b is detected at much lower levels in kidney, liver, brain and testis and is not detected in heart or skeletal muscle.		

— VALIDATION IMAGES —

Sample: Lane 1: Kidney (Mouse) Lysate at 40 ug
 Lane 2: Cerebellum (Mouse) Lysate at 40 ug
 Lane 3: Cerebrum (Mouse) Lysate at 40 ug
 Lane 4: Cerebellum (Rat) Lysate at 40 ug
 Primary: Anti-PANK1 (bs-8337R) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 60/50 kD
 Observed band size: 60/50 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PANK1) Polyclonal Antibody, Unconjugated (bs-8337R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.