

**bs-1417R****[ Primary Antibody ]**

**BioSS**  
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**NFAT2 Rabbit pAb****— DATASHEET —**


|   |                      |  |
|---|----------------------|--|
| <b>Host:</b> Rabbit   | <b>Isotype:</b> IgG  | <b>Applications:</b> IHC-P (1:100-500)<br>IHC-F (1:100-500)<br>IF (1:100-500)<br>Flow-Cyt (1ug/test) |
| <b>Clonality:</b> Polyclonal  |                      |  |
| <b>GeneID:</b> 4772   | <b>SWISS:</b> O95644 |  |
| <b>Target:</b> NFAT2  |                      |  |
| <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human NFATc1: 601-716/716.  |                      | <b>Reactivity:</b> Human, Rat<br>(predicted: Mouse)  |
| <b>Purification:</b> affinity purified by Protein A   |                      |  |
| <b>Concentration:</b> 1mg/ml  |                      | <b>Predicted MW.:</b> 101 kDa  |
| <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.<br>Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.  |                      | <b>Subcellular Location:</b> Cytoplasm ,Nucleus  |
| <b>Background:</b> The product of this gene is a component of the nuclear factor of activated T cells DNA-binding transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation, and an inducible nuclear component. Proteins belonging to this family of transcription factors play a central role in inducible gene transcription during immune response. The product of this gene is an inducible nuclear component. It functions as a major molecular target for the immunosuppressive drugs such as cyclosporin A. Multiple alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. Different isoforms of this protein may regulate inducible expression of different cytokine genes. [provided by RefSeq, Jul 2013] |                      |  |

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NFAT2) Polyclonal Antibody, Unconjugated (bs-1417R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: Molt-4. Primary Antibody (green line): Rabbit Anti-NFAT2 antibody (bs-1417R) Dilution: 1μg / 10<sup>6</sup> cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1μg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control: Jurkat. Primary Antibody (green line): Rabbit Anti-NFAT2 antibody (bs-1417R) Dilution: 1ug/Test; Secondary Antibody : Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

**— SELECTED CITATIONS —**

- **[IF=14.976]** Qinyu Ma. et al. Small extracellular vesicles deliver osteolytic effectors and mediate cancer - induced

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

- osteolysis in bone metastatic niche. J Extracell Vesicles. 2021 Feb;10(4):e12068 WB ;Mouse. 33659051
- **[IF=8.063]** Zhang Y et al. ASIC1a induces synovial inflammation via the Ca<sup>2+</sup>/NFATc3/RANTES pathway. Theranostics. 2020 Jan 1;10(1):247-264. IF ;Rat. 31903118
  - **[IF=8.109]** Ling Wu. et al. Calcium Channel Blocker Nifedipine Suppresses Colorectal Cancer Progression and Immune Escape by Preventing NFAT2 Nuclear Translocation. Cell Rep. 2020 Oct;33:108327 IF,IHC ;Human. 33113363
  - **[IF=5.6]** Yu Bai. et al. LMCD1 is involved in tubulointerstitial inflammation in the early phase of renal fibrosis by promoting NFATc1-mediated NLRP3 activation. INT IMMUNOPHARMACOL. 2023 Aug;121:110362 IF,ICC ;Mouse,Human. 37311356
  - **[IF=5.039]** Zhang B et al. miR-137 and its target T-type CaV 3.1 channel modulate dedifferentiation and proliferation of cerebrovascular smooth muscle cells in simulated microgravity rats by regulating calcineurin/NFAT pathway. Cell Proliferation. 2020;00:e12774. WB ;Rat. doi:10.1111/cpr.12774