

bs-16214R**[Primary Antibody]****GAL3ST2 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
Clonality: Polyclonal		
GeneID: 64090	SWISS: Q9H3Q3	
Target: GAL3ST2		Reactivity: Rat (predicted: Human, Mouse, Rabbit, Pig, Cow, Chicken, Dog, Horse)
Immunogen: KLH conjugated synthetic peptide derived from human GAL3ST2: 1-100/398.		
Purification: affinity purified by Protein A		Predicted MW.: 46 kDa
Concentration: 1mg/ml		Subcellular Location: Cytoplasm
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: This gene encodes a member of the galactose-3-O-sulfotransferase protein family. The product of this gene catalyzes sulfonation by transferring a sulfate group to the hydroxyl at C-3 of nonreducing beta-galactosyl residues, and it can act on both type 1 and type 2 (Galbeta 1-3/1-4GlcNAc-R) oligosaccharides with similar efficiencies, and on core 1 glycans. This enzyme has been implicated in tumor metastasis processes. This gene is different from the GAL3ST3 gene located on chromosome 11, which has also been referred to as GAL3ST2 and encodes a related enzyme with distinct tissue distribution and substrate specificities, compared to galactose-3-O-sulfotransferase 2. [provided by RefSeq, Jul 2008]		

— VALIDATION IMAGES —

Tissue/cell: Rat spleen tissue; 4%
Paraformaldehyde-fixed and paraffin-
embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block
endogenous peroxidase by 3% Hydrogen
peroxide for 30min; Blocking buffer (normal goat
serum, C-0005) at 37°C for 20 min; Incubation:
Anti-GAL3ST2 Polyclonal Antibody,
Unconjugated(bs-16214R) 1:200, overnight at
4°C, followed by conjugation to the secondary
antibody(SP-0023) and DAB(C-0010) staining