

bs-16221R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

GALNT5 Rabbit pAb**— DATASHEET —**

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 11227</p> <p>Target: GALNT5</p> <p>Immunogen: KLH conjugated synthetic peptide derived from human GALNT5: 501-600/940.</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: The UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase (GalNAc-T) family of enzymes are substrate-specific proteins that catalyze the transfer of GalNAc (N-acetylgalactosamine) to serine and threonine residues onto various proteins, thereby initiating mucin-type O-linked glycosylation in the Golgi apparatus. GalNAc-T5 (Polypeptide N-acetylgalactosaminyltransferase 5), also known as UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase 5, is a 940 amino acid protein that displays enzymatic activity toward EA2 peptide substrate with weaker activity toward Muc2 or Muc 1b substrates. Its N-terminal domain is involved in substrate binding and manganese coordination, while the C-terminal domain is involved in UDP-Gal binding and catalytic reaction. EXT2 directly interacts with GalNAc-T5, suggesting that these proteins may corroborate in glycosaminoglycan synthesis.</p>	<p>Isotype: IgG</p> <p>SWISS: Q7Z7M9</p>	<p>Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000)</p> <p>Reactivity: (predicted: Human, Mouse)</p> <p>Predicted MW.: 106 kDa</p> <p>Subcellular Location: Cell membrane</p>
--	--	--