bs-13187R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn

techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

FMO5 Rabbit pAb

GenelD: 2330 **SWISS:** P49326

Target: FMO5

Immunogen: KLH conjugated synthetic peptide derived from human FMO5:

251-350/533.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The Flavin containing monooxygenase family consists of five gene products, FMO1-5, that are major enzymatic oxidants involved in the metabolism of various therapeutics. Localizing to microsomal and endoplasmic reticulum membranes, FMO5 (flavin containing monooxygenase 5), also known as dimethylaniline monooxygenase [N-oxide-forming] 5, hepatic flavin-containing monooxygenase 5 or dimethylaniline oxidase 5, is a 533 amino acid protein belonging to the FMO family. Expressed in adult and fetal liver, FMO5 is unlike other FMO family members because it does not function as a drug-metabolizing enzyme. FMO5 binds FAD as a cofactor and is encoded by a gene located on human chromosome

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Human, Mouse

(predicted: Rat, Pig, Cow,


Dog)

Predicted 60 kDa


Subcellular Location: Cell membrane ,Cytoplasm

VALIDATION IMAGES

1q21.1.



Protein: liver(mouse)lyates at 40ug; Primary: Anti-FMO5 (bs-13187R) at 1:300; Secondary: HRP conjugated Goat-Anti-Rabbit IgG(bse-0295G-HRP) at 1: 5000; ECL excitated the fluorescence; Predicted band size: 60 kD Observed band size



Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-FMO5 Polyclonal Antibody. Unconjugated(bs-13187R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining