
5HT2C Receptor Rabbit pAb

Catalog Number: bs-2959R

Target Protein: 5HT2C Receptor

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse (predicted:Rat, Pig, Cow, Dog, Horse)

Predicted MW: 52 kDa

Entrez Gene: 3358

Swiss Prot: P28335

Source: KLH conjugated synthetic peptide derived from human 5HTR2C: 311-410/458.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Serotonin (5-hydroxytryptamine, 5-HT), a neurotransmitter, elicits a wide array of physiological effects by binding to several receptor subtypes, including the 5-HT2 family of seven-transmembrane-spanning, G-protein-coupled receptors, which activate phospholipase C and D signaling pathways. This gene encodes the 2C subtype of serotonin receptor and its mRNA is subject to multiple RNA editing events, where genomically encoded adenosine residues are converted to inosines. RNA editing is predicted to alter amino acids within the second intracellular loop of the 5-HT2C receptor and generate receptor isoforms that differ in their ability to interact with G proteins and the activation of phospholipase C and D signaling cascades, thus modulating serotonergic neurotransmission in the central nervous system. Studies in humans have reported abnormalities in patterns of 5-HT2C editing in depressed suicide victims. [provided by RefSeq, Jul 2008].

VALIDATION IMAGES



Sample: Lane 1: Human SH-SY5Y cell lysates Lane 2: Human A431 cell lysates Primary: Anti-5HT2C Receptor (bs-2959R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52 kDa Observed band size: 59 kDa



Paraformaldehyde-fixed, paraffin embedded (mouse cerebellum tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (5HTR2C) Polyclonal Antibody, Unconjugated (bs-2959R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

PRODUCT SPECIFIC PUBLICATIONS

[IF=14.971] Kyritsi K et al. Modulation of the TPH 1/MAO - A/5 HT/5 HTR 2A/2B/2C Axis Regulates Biliary Proliferation and Liver Fibrosis During Cholestasis. Hepatology.2019 Jul 25. ICC,IHC ; Rat&Mouse . doi:10.1002/hep.30880

[IF=4.546] Quancheng Liu. et al. FumDSB Can Reduce the Toxic Effects of Fumonisin B1 by Regulating Several Brain-Gut Peptides in Both the Hypothalamus and Jejunum of Growing Pigs. Toxins. 2021 Dec;13(12):874 WB,IHC ; Pig . 34941712