

bs-6305R**[Primary Antibody]**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Ryanodine Receptor Rabbit pAb**— DATASHEET —**

Host: Rabbit Clonality: Polyclonal GeneID: 6261 Target: Ryanodine Receptor Immunogen: KLH conjugated synthetic peptide derived from human Ryanodine Receptor: 4701-4800/5038. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The Ryanodine Receptor (RyR) is the channel responsible for calcium release from muscle cell Sarcoplasmic Reticulum (SR) and also plays a role in calcium regulation in non-muscle cells. The RyR exists as a homotetramer and is predicted to have a short cytoplasmic C-terminus and 4-10 transmembrane domains. The remainder of the protein, termed the "foot" region, is located in the cytoplasm between the transverse tubule and the SR. Mammalian RyR isoforms are the product of three different genes: RyR-1 is expressed predominantly in skeletal muscle and areas of the brain; RyR-2 is expressed predominantly in heart muscle but also found in the stomach, endothelial cells and diffuse areas of the brain; and RyR-3 is found in smooth muscle and the brain (striatum, thalamus and hippocampus). In non-mammalian vertebrates, the RyR isoforms are termed alpha, beta and cardiac which correlate loosely to the mammalian RyR-1, RyR-3 and RyR-2 isoforms respectively.	Isotype: IgG SWISS: P21817 Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Mouse (predicted: Human, Rat, Rabbit, Pig, Cow, Dog) Predicted MW.: 566 kDa Subcellular Location: Cell membrane ,Cytoplasm
--	---

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (mouse cerebellum tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Ryanodine Receptor) Polyclonal Antibody, Unconjugated (bs-6305R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=2.466]** Gerburg Keilhoff. et al. The Ryanodine receptor stabilizer S107 fails to support motor neuronal neuritogenesis in vitro. Tissue Cell. 2021 Dec;73:101625 IF ;Mouse. 34419737

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.