bs-10802R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

TNF alpha Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 7124 **SWISS:** P01375

Target: TNF alpha

Immunogen: Recombinant human TNF alpha protein: 77-233/233.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. This cytokine is mainly secreted by macrophages. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, psoriasis, rheumatoid arthritis ankylosing spondylitis, tuberculosis, autosomal dominant polycystic kidney disease, and cancer. Mutations in this gene affect susceptibility to cerebral malaria, septic shock, and Alzheimer disease. Knockout studies in mice also suggested the neuroprotective function of this cytokine. [provided by RefSeq, Aug 2020]


Applications: Flow-Cyt (1ug/Test) ICC/IF (1:100-500)

Reactivity: Human, Mouse


Predicted MW.: 17/26 kDa

Subcellular Secreted ,Cell membrane


VALIDATION IMAGES



4% Paraformaldehyde-fixed THP-1(LPS treated (100 ng/ml, 16 h) and Brefeldin A treated (5 μg/ml, 4 h)) (H) cell; Antibody incubation with (TNF alpha) polyclonal Antibody, unconjugated (bs-10802R) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-60295G-BF488) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.



Blank control:Raw264.7. Primary Antibody (green line): Rabbit Anti-TNF alpha antibody (bs-10802R) Dilution: 1ug/Test; Secondary Antibody: Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test, Protocol The cells were treated with LPS (1 ug/mL, 18 hr/6 hr) and Brefeldin A (300 ng/mL, last 3 hr of stimulation). The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block nonspecific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control:THP-1. Primary Antibody (green line): Rabbit Anti-TNF alpha antibody (bs-10802R) Dilution: 1ug/Test; Secondary Antibody: Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test, Protocol The cells were treated with TPA (80 nM, overnight) and then treated with LPS (1 ug/mL, 18 hr/6 hr) and Brefeldin A (300 $\,$ ng/mL, last 3 hr of stimulation) .The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block nonspecific protein-protein interactions for 30 min at room temperature . Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

- [IF=21.3] Gu Xiaoxue. et al. Bismuth-Nanosheet-Armed Pristine Silk Nanofiber Dressing for Multimodal Pathogenic Bacteria Eradication and Infected Wound Healing. ADV FIBER MATER. 2025 Jun;:1-16 IHC; Mouse. 10.1007/s42765-025-00568-z
- [IF=19] Qianqian Liu. et al. Montmorillonite Modulates Manganese d-Band Center to Enhance Cascade Reaction Activity Against Inflammatory Bowel Disease. ADV FUNCT MATER. 2025 May;:2502389 |F; Mouse. 10.1002/adfm.202502389
- [IF=17.521] Huan Lei. et al. A Combination Therapy Using Electrical Stimulation and Adaptive, Conductive Hydrogels Loaded with Self-Assembled Nanogels Incorporating Short Interfering RNA Promotes the Repair of Diabetic Chronic Wounds. Advanced Science. 2022 Sep;;2201425 IF; Rat. 36064844
- [IF=17.694] Wang, Yi. et al. In-situ growth of robust superlubricated nano-skin on electrospun nanofibers for post-operative adhesion prevention. NAT COMMUN. 2022 Aug;13(1):1-12 IF; Rat. 36030284
- [IF=16.744] Lubin Zhou. et al. A self-pumping dressing with in situ modification of non-woven fabric for promoting diabetic wound healing. CHEM ENG J. 2022 Dec;:141108 IHC; Rat. 10.1016/j.cej.2022.141108