— DATASHEET —

[Primary Antibody]


phospho-DNA PKcs (Thr2638) Rabbit pAb




www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500) IF (1:100-500)
GenelD: 5591	SWISS: P78527	IF (1.100-500)
Target: DNA PKcs (Thr2638)		Reactivity: Mouse (predicted: Human, Rat)
	thesised phosphopeptide derived from h e phosphorylation site of Thr2638: RA(p-1	
Purification: affinity purified by Protein A		Predicted MW.: ^{454 kDa}
Concentration: 1mg/ml		MW.: 15 TKBU
Storage: Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		PBS, Subcellular Nucleus
protein kinase (DNA- heterodimer protein recombination. The	ne catalytic subunit of the DNA-depender PK). It functions with the Ku70/Ku80 in DNA double strand break repair and protein encoded is a member of the PI3/I led by RefSeq, Jul 2010]	

– VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-DNA PKcs (Thr2638)) Polyclonal Antibody, Unconjugated (bs-20347R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-DNA PKcs (Thr2638)) Polyclonal Antibody, Unconjugated (bs-20347R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.