

bs-18036R**[Primary Antibody]****ornithine aminotransferase Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 4942	SWISS: P04181	IHC-F (1:100-500)
Target: ornithine aminotransferase		IF (1:100-500)
Immunogen: KLH conjugated synthetic peptide derived from human ornithine aminotransferase: 351-439/439.		Reactivity: Human, Mouse, Rat (predicted: Pig, Cow, Chicken, Horse)
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Predicted MW.: 45 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cytoplasm
Background: OAT (ornithine aminotransferase (mitochondrial), ornithine-oxo-acid aminotransferase) is a 439 amino acid protein encoded by the human gene OAT. OAT belongs to the class III pyridoxal-phosphate-dependent aminotransferase family and is usually found as a homotetramer in the mitochondrion matrix. OAT catalyzes the major catalytic reaction for ornithine. Ornithinemia, presumably due to deficiency of ornithine ketoacid aminotransferase (OAT) has been found in patients with gyrate atrophy of the choroid and retina. The clinical history of gyrate atrophy is usually night blindness that begins in late childhood, accompanied by sharply demarcated circular areas of chorioretinal atrophy. During the second and third decades the areas of atrophy enlarge. The hepatic cleavage product, hepatic OAT, is formed by cleaving a 25 amino acid transit peptide from the N-terminus of the OAT precursor. The renal form is produced by cleaving a 35 amino acid transit peptide from the N-terminus.		

— VALIDATION IMAGES —

Sample: A431(Human) Cell Lysate at 30 ug Small intestine (Mouse) Lysate at 40 ug Primary: Anti-ornithine aminotransferase (bs-18036R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 45 kD Observed band size: 45 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ornithine aminotransferase) Polyclonal Antibody, Unconjugated (bs-18036R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.