bs-11666R

[Primary Antibody]

PPP1R10 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

— DATASHEET ————		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GenelD: 5514	SWISS: Q96QC0	IF (1:100-500)
Target: PPP1R10		ICC/IF (1:25)
Immunogen: KLH conjugated synthetic peptide derived from human PPP1R10: 151-250/940.		P1R10: Reactivity: Human (predicted: Mouse, Rat, Pig, Sheep, Cow, Dog)
Purification: affinity purified by I	Protein A	
Concentration: 1mg/ml		Predicted
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.		MW.: ^{99 kDa}
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Nucleus
Background: Eukaryotic protein phosphorylation and dephosphorylation on serine and threonine residues regulates numerous cell functions, including division, homeostasis and apoptosis. A group of proteins that play a major role in this process are the serine/threonine protein phosphatases. Protein phosphatase (PP) holoenzyme is a trimeric complex that contains a regulatory subunit, a variable subunit and a catalytic subunit. Families of PP catalytic subunits include PP1, PP2A, PP2B, PP2C, PPX and PP5. Regulatory subunits include nuclear inhibitor of PP1 (NIPP1), PP1 nuclear targeting subunit (PNUTS), PP2A-A, PP2A-B, PP2A-B56, PP2A-C, PP2B-B and PR48. PNUTS, also designated CAT53 or FB19, is encoded by the gene PPP1R10. PNUTS acts as an inhibitor for the phosphatase activity of PP1 Alpha and PP1 Gamma. It is a nuclear protein primarily detected in nucleoplasmic bodies and within nucleoli. PNUTS expression levels are highest in brain, heart, lung, placenta, liver, kidney, pancreas and skeletal muscle.		n on tions, proteins ne me is a ible punits ubunits ubunits ing B-B and y the tase n leoli. lacenta,

- VALIDATION IMAGES -



Sample: Lane 1: Human MCF-7 cell lysates Lane 2: Human HeLa cell lysates Lane 3: Human HepG2 cell lysates Primary: Anti-PPP1R10 (bs-11666R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 99 kDa Observed band size: 115 kDa



Tissue/cell: Human kidney tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-ppp1R10 Polyclonal Antibody, Unconjugated(bs-11666R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (PPP1R10) polyclonal Antibody, Unconjugated (bs-11666R) 1:25, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.