

**bs-16236R**

**[ Primary Antibody ]**

## GDA/GDC Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

### — DATASHEET —

<p><b>Host:</b> Rabbit</p> <p><b>Clonality:</b> Polyclonal</p> <p><b>GeneID:</b> 9615</p> <p><b>Target:</b> GDA/GDC</p> <p><b>Immunogen:</b> KLH conjugated synthetic peptide derived from human GDA: 21-120/454.</p> <p><b>Purification:</b> affinity purified by Protein A</p> <p><b>Concentration:</b> 1mg/ml</p> <p><b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p><b>Background:</b> This gene encodes an enzyme responsible for the hydrolytic deamination of guanine. Studies in rat ortholog suggest this gene plays a role in microtubule assembly. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2011]</p>	<p><b>Isotype:</b> IgG</p> <p><b>SWISS:</b> Q9Y2T3</p> <p><b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>ICC/IF</b> (1:100-500) <b>ELISA</b> (1:5000-10000)</p> <p><b>Reactivity:</b> (predicted: Human, Mouse, Rat, Rabbit, Cow, Dog, Horse)</p> <p><b>Predicted MW.:</b> 51 kDa</p> <p><b>Subcellular Location:</b> Secreted ,Extracellular matrix ,Cytoplasm</p>
---	---