

bs-16249R

[Primary Antibody]

GLIS3 Rabbit pAb

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000) Reactivity: Human, Rat, Sheep Predicted MW.: 84 kDa Subcellular Location: Nucleus
Clonality: Polyclonal		
GeneID: 169792	SWISS: Q8NEA6	
Target: GLIS3		
Immunogen: KLH conjugated synthetic peptide derived from human GLIS3: 351-450/775.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: This gene is a member of the GLI-similar zinc finger protein family and encodes a nuclear protein with five C2H2-type zinc finger domains. This protein functions as both a repressor and activator of transcription and is specifically involved in the development of pancreatic beta cells, the thyroid, eye, liver and kidney. Mutations in this gene have been associated with neonatal diabetes and congenital hypothyroidism (NDH). Alternatively spliced variants that encode different protein isoforms have been described but the full-length nature of only two have been determined. [provided by RefSeq, Jul 2008]		


— VALIDATION IMAGES —



Sample: Lane 1: Human U251 cell lysates Lane 2: Human U87MG cell lysates Lane 3: Human A549 cell lysates Lane 4: Human 293T cell lysates Lane 5: Human MDA-MB-231 cell lysates Primary: Anti-GLIS3 (bs-16249R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 84 kDa Observed band size: 84 kDa



Tissue/cell: human kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-GLIS3 Polyclonal Antibody, Unconjugated(bs-16249R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-GLIS3 Polyclonal Antibody, Unconjugated(bs-16249R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

— SELECTED CITATIONS —

- **[IF=3.252]** Zhuohang Liu. et al. GLIS Family Zinc Finger 3 Promoting Cell Malignant Behaviors and NF-κB Signaling in Glioma. Brain Res. 2021 Aug;;147623 IHC ;Human. 34403660
- **[IF=2.7]** Bing Wang. et al. Co-inhibition of adenosine 2b receptor and programmed death-ligand 1 promotes the recruitment and cytotoxicity of natural killer cells in oral squamous cell carcinoma. PEERJ. 2023 Aug;11:e15922 IHC ;Human. 37663280