bs-19704R

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

R3HDM2 Rabbit pAb

GenelD: 22864 **SWISS:** Q9Y2K5

Target: R3HDM2

Immunogen: KLH conjugated synthetic peptide derived from human R3HDM2:

701-800/976.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The R3H motif is characterised by the presence of an invariant

arginine residue and a highly conserved histidine residue, which are separated by three residues. The function of the domain is predicted to be binding ssDNA. There are two named isoforms of

R3HDM2.

Applications: IHC-P (1:100-500)


IHC-F (1:100-500) IF (1:100-500)

Reactivity: Human (predicted: Mouse)


Predicted MW.: 107 kD

Subcellular Nucleus

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (R3HDM2) Polyclonal Antibody, Unconjugated (bs-19704R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (R3HDM2) Polyclonal Antibody, Unconjugated (bs-19704R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.