

bs-8865R**[Primary Antibody]****CD28 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 940 Target: CD28 Immunogen: KLH conjugated synthetic peptide derived from human CD28: 19-120/220. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The protein encoded by this gene is essential for T-cell proliferation and survival, cytokine production, and T-helper type-2 development. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2011]	Isotype: IgG SWISS: P10747	Applications: WB (1:500-2000) ELISA (1:5000-10000) Reactivity: Mouse (predicted: Human, Rat) Predicted MW.: 25 kDa Subcellular Location: Cell membrane
--	---	--

— SELECTED CITATIONS —

- **[IF=3.549]** Jingjing Li. et al. Circular RNA hsa_circ_0068252 functions in cisplatin resistance and immune response via miR-1304-5p/PD-L1 axis in non-small cell lung cancer. CHEMOTHERAPY. ;: Other ;Human. 35649347
- **[IF=3.057]** Song et al. Forsythoside A Inhibits BVDV Replication via TRAF2-Dependent CD28-4-1BB Signaling in Bovine PBMCs. (2016) PLoS.On. 11:e0162791 WB ;COW. 27617959
- **[IF=3.098]** Junli Zhang. et al. KCNQ1OT1 contributes to sorafenib resistance and programmed death-ligand-1-mediated immune escape via sponging miR-506 in hepatocellular carcinoma cells. Int J Mol Med. 2020 Nov;46(5):1794-1804 ELISA ;Human. 33000204