### bs-6811R

## [ Primary Antibody ]

# Bioss ANTIBODIES

## www.bioss.com.cn sales@bioss.com.cn

Applications: IHC-P (1:100-500)

Reactivity: Human, Mouse

**Predicted** 

MW.:

Subcellular Nucleus

techsupport@bioss.com.cn 400-901-9800

IHC-F (1:100-500)

(predicted: Rat, Sheep, Cow, Dog, Horse)

**IF** (1:100-500)

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

MUM1 Rabbit pAb

GenelD: 84939 SWISS: Q2TAK8

Target: MUM1

**Immunogen:** KLH conjugated synthetic peptide derived from human

MUM1/FLJ14868: 401-500/710.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated


freeze/thaw cycles.

**Background:** Enables nucleosome binding activity. Involved in DNA repair and


chromatin organization. Located in cytosol and nucleoplasm.

[provided by Alliance of Genome Resources, Jun 2025]

#### VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MUM1) Polyclonal Antibody, Unconjugated (bs-6811R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse pancreas tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MUM1) Polyclonal Antibody, Unconjugated (bs-6811R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.