

**bs-6811R****[ Primary Antibody ]****Bioss**  
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**MUM1 Rabbit pAb****— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)  <b>Reactivity:</b> Human, Mouse (predicted: Rat, Sheep, Cow, Dog, Horse)  <b>Predicted MW.:</b> 79 kDa  <b>Subcellular Location:</b> Cell membrane ,Cytoplasm
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 84939	<b>SWISS:</b> Q2TAK8	
<b>Target:</b> MUM1		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human MUM1/FLJ14868: 401-500/710.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> Munc 13s are homologues of the C. elegans unc-13 gene product. Three brain specific isoforms are expressed in rat: Munc 13-1, -13-2, and -13-3. Munc 13-1 is a peripheral membrane protein that is enriched in synaptosomes and localized to plasma membranes but absent from synaptic vesicles. It has multiple regulatory domains (C1 and C2 domains) that may mediate diacylglycerol, phospholipid, and calcium binding. The C terminus of Munc 13-1 interacts directly with a putative coiled coil domain in the N-terminal part of syntaxin. It has been shown that Munc 13 is involved in priming synaptic vesicles. These activated vesicles can be released immediately on Ca <sup>2+</sup> influx.		

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MUM1) Polyclonal Antibody, Unconjugated (bs-6811R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse pancreas tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MUM1) Polyclonal Antibody, Unconjugated (bs-6811R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.