

bs-18631R**[Primary Antibody]****MAGOH/Mago nashi homolog 2 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Mouse, Rat (predicted: Human, Cow, Dog) Predicted MW.: 17 kDa Subcellular Location: Nucleus
Clonality: Polyclonal		
GeneID: 55110	SWISS: Q96A72	
Target: MAGOH/Mago nashi homolog 2		
Immunogen: KLH conjugated synthetic peptide derived from human MAGOH/Mago nashi homolog 2: 51-148/148.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: MAGOHB (Mago Homolog B, Exon Junction Complex Subunit) is a Protein Coding gene. Diseases associated with MAGOHB include Metaphyseal Chondrodysplasia, Schmid Type and Chromosome 22Q11.2 Deletion Syndrome, Distal. Among its related pathways are Cleavage of Growing Transcript in the Termination Region and Transport of Mature Transcript to Cytoplasm. An important paralog of this gene is MAGOH.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MAGOH) Polyclonal Antibody, Unconjugated (bs-18631R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MAGOH) Polyclonal Antibody, Unconjugated (bs-18631R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.