

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

STK4 Rabbit pAb

Catalog Number: bs-55209R

Target Protein: STK4
Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Human, Mouse, Rat

Predicted MW: 56 kDa Entrez Gene: 6789 Swiss Prot: Q13043

Source: Recombinant human STK4: 303-487/487.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is

stable for at least two weeks at 2-4°C.

Background: The protein encoded by this gene is a cytoplasmic kinase that is structurally similar to the

yeast Ste20p kinase, which acts upstream of the stress-induced mitogen-activated protein


kinase cascade. The encoded protein can phosphorylate myelin basic protein and undergoes autophosphorylation. A caspase-cleaved fragment of the encoded protein has been shown to be capable of phosphorylating histone H2B. The particular phosphorylation catalyzed by this protein has been correlated with apoptosis, and it's possible that this

catalyzed by this protein has been correlated with apoptosis, and it's possible that this

protein induces the chromatin condensation observed in this process. [provided by RefSeq,

Jul 2008]

VALIDATION IMAGES



Sample: Lane 1: Cerebrum (Mouse) Lysate at 40 ug Lane 2: Testis (Mouse) Lysate at 40 ug Lane 3: Spleen (Mouse) Lysate at 40 ug Lane 4: Lymph node (Mouse) Lysate at 40 ug Lane 5: NIH/3T3 (Mouse) Cell Lysate at 30 ug Lane 6: Cerebrum (Rat) Lysate at 40 ug Lane 7: Testis (Rat) Lysate at 40 ug Lane 8: Hela (Human) Cell Lysate at 30 ug Lane 9: Jurkat (Human) Cell Lysate at 30 ug Lane 10: HL60 (Human) Cell Lysate at 30 ug Lane 11: Molt-4 (Human) Cell Lysate at 30 ug Primary: Anti-STK4 (bs-55209R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 60 kD Observed band size: 58 kD