bs-55209R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

STK4 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 6789 **SWISS:** Q13043

Target: STK4

Immunogen: Recombinant human STK4: 303-487/487.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the

antibody is stable for at least two weeks at 2-4°C.

Background: The protein encoded by this gene is a cytoplasmic kinase that is

structurally similar to the yeast Ste20p kinase, which acts upstream of the stress-induced mitogen-activated protein kinase cascade. The encoded protein can phosphorylate myelin basic protein and undergoes autophosphorylation. A caspase-cleaved fragment of the encoded protein has been shown to be capable of phosphorylating histone H2B. The particular phosphorylation catalyzed by this protein has been correlated with apoptosis, and it's possible that this protein induces the chromatin condensation

observed in this process. [provided by RefSeq, Jul 2008]

Applications: WB (1:500-2000)

ICC/IF (1:50 - 1:100) **ELISA** (1:5000-10000) **IP** (0.5μg-4μg antibody for 200µg-400µg extracts of


whole cells)

Reactivity: Human, Mouse, Rat

Predicted MW.: 56 kDa

Subcellular Cytoplasm ,Nucleus

VALIDATION IMAGES



Sample: Lane 1: Cerebrum (Mouse) Lysate at 40 ug Lane 2: Testis (Mouse) Lysate at 40 ug Lane 3: Spleen (Mouse) Lysate at 40 ug Lane 4: Lymph node (Mouse) Lysate at 40 ug Lane 5: NIH/3T3 (Mouse) Cell Lysate at 30 ug Lane 6: Cerebrum (Rat) Lysate at 40 ug Lane 7: Testis (Rat) Lysate at 40 ug Lane 8: Hela (Human) Cell Lysate at 30 ug Lane 9: Jurkat (Human) Cell Lysate at 30 ug Lane 10: HL60 (Human) Cell Lysate at 30 ug Lane 11: Molt-4 (Human) Cell Lysate at 30 ug Primary: Anti-STK4 (bs-55209R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 60 kD Observed band size: 58 kD