## bs-11968R

## [ Primary Antibody ]

# BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# Semaphorin 3D Rabbit pAb

- DATASHEET -

**Host:** Rabbit **Isotype:** IgG

Clonality: Polyclonal

**GenelD:** 223117 **SWISS:** 095025

Target: Semaphorin 3D

**Immunogen:** KLH conjugated synthetic peptide derived from human

Semaphorin 3D: 351-450/777.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** Semaphorins are a family of cell surface and secreted proteins that

are conserved from insects to humans. Members of this family of proteins are approximately 750 amino acids in length (including signal sequences) and are defined by a conserved extracellular "semaphorin" domain of approximately 500 amino acids containing 14-16 cysteines, blocks of conserved sequences and no obvious repeats. Secreted and cell-bound semaphorins chemically attract and repel the growth of neural axons, guiding the development of intricate networks of neural tissue. SEMA3D (semaphorin-3D), also known as coll-2 or Sema-Z2, is a 777 amino acid secreted protein that belongs to the semaphorin family and participates in axon-axon interactions and neural crest activity. SEMA3D binds neuropilin and is able to induce the collapse and

paralysis of neuronal growth cones, as well as act to repel specific neuronal populations. SEMA3D contains one immunoglobulin-like (Ig-like) domain, one PSI domain and one semaphorin domain.

Applications: IHC-P (1:100-500)

**IHC-F** (1:100-500) **IF** (1:100-500)


Reactivity: Mouse, Rat

(predicted: Human, Rabbit, Pig, Chicken, Dog, Horse)


Predicted MW.: 85 kDa

Subcellular Secreted

#### VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by microwave in sodium citrate buffer (pH6.0); Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes; Blocking buffer (3% BSA) at RT for 30min; Antibody incubation with (Semaphorin 3D) Polyclonal Antibody, Unconjugated (bs-11968R) at 1:400 overnight at 4°C, followed by conjugation to the secondary antibody (labeled with HRP) and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by microwave in sodium citrate buffer (pH6.0); Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes; Blocking buffer (3% BSA) at RT for 30min; Antibody incubation with (Semaphorin 3D) Polyclonal Antibody, Unconjugated (bs-11968R) at 1:400 overnight at 4°C, followed by conjugation to the secondary antibody (labeled with HRP) and DAB staining.

### - SELECTED CITATIONS -

• [IF=3.26] Liu, Xin-Qi, et al. "Regulation of neuroendocrine cells and neuron factors in the ovary by zinc oxide nanoparticles." Toxicology Letters (2016). IHC; Chicken. 27215404