

bs-11972R**[Primary Antibody]****RAMP3 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) ELISA (1:5000-10000)
Clonality: Polyclonal		
GeneID: 10268	SWISS: O60896	
Target: RAMP3		
Immunogen: KLH conjugated synthetic peptide derived from human RAMP3: 51-148/148.		
Purification: affinity purified by Protein A		Reactivity: Human, Mouse
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 14 kDa
Background: Calcitonin receptor like receptor activity modifying protein 3; CRLR activity modifying protein 3; CRLR activity-modifying protein 3; OTTHUMP00000159549; OTTHUMP00000214279; OTTHUMP00000214280; Receptor (calcitonin) activity modifying protein 3; Receptor (G protein-coupled) activity modifying protein 3; Receptor activity modifying protein 3.		Subcellular Location: Cell membrane

— VALIDATION IMAGES —

25 ug total protein per lane of various lysates (see on figure) probed with RAMP3 polyclonal antibody, unconjugated (bs-11972R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.

— SELECTED CITATIONS —

- **[IF=3.3]** Wang Qinghao. et al. Eosinophil-Associated Genes are Potential Biomarkers for Hepatocellular Carcinoma Prognosis. J CANCER. 2024 Sep;15(17):5605-5621 IHC ;Human. 39308686
- **[IF=1.86]** Qiao, Xi, et al. "Intermedin is upregulated and attenuates renal fibrosis by inhibition of oxidative stress in rats with unilateral ureteral obstruction."Nephrology (2015). WB ;="Rat". 26014968