bs-11972R

[Primary Antibody]

www.bioss.com.cn

RAMP3 Rabbit pAb

sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 10268 **SWISS:** 060896

Target: RAMP3

Immunogen: KLH conjugated synthetic peptide derived from human RAMP3:

51-148/148. < Extracellular >

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Calcitonin receptor like receptor activity modifying protein 3; CRLR

activity modifying protein 3; CRLR activity-modifying protein 3;

OTTHUMP00000159549; OTTHUMP00000214279;

OTTHUMP00000214280; Receptor (calcitonin) activity modifying protein 3; Receptor (G protein-coupled) activity modifying protein

3; Receptor activity modifying protein 3.

Applications: WB (1:500-2000)

ELISA (1:5000-10000)


Reactivity: Human, Mouse

Predicted 14 kDa

MW.:

Subcellular Cell membrane

VALIDATION IMAGES -



25 ug total protein per lane of various lysates (see on figure) probed with RAMP3 polyclonal antibody, unconjugated (bs-11972R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at rt for 60 min

— SELECTED CITATIONS –

- [IF=3.3] Wang Qinghao. et al. Eosinophil-Associated Genes are Potential Biomarkers for Hepatocellular Carcinoma Prognosis. J CANCER. 2024 Sep;15(17):5605-5621 IHC; Human. 39308686
- [IF=1.86] Qiao, Xi, et al. "Intermedin is upregulated and attenuates renal fibrosis by inhibition of oxidative stress in rats with unilateral ureteral obstruction." Nephrology (2015). WB; Rat. 26014968