
FTS Rabbit pAb

Catalog Number: bs-13223R

Target Protein: FTS

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Rat (predicted:Human, Mouse, Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse)

Predicted MW: 33 kDa

Subcellular: Cell membrane ,Cytoplasm

Locations:

Entrez Gene: 64400

Swiss Prot: Q9H8T0

Source: KLH conjugated synthetic peptide derived from human FTS/AKTIP: 201-292/292.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Fused toes protein homolog (FTS), also known as AKT-interacting protein (AKTIP) and Ft1, is a 292 amino acid protein that localizes to the cytoplasm and the cell membrane. A member of the ubiquitin-conjugating enzyme family, FTS binds directly to AKT1 to regulate apoptosis in a cell population. AKT1 is a protein that plays a critical role in a number of cellular responses, such as cell growth, protein synthesis, and antiapoptotic signaling. The interaction of FTS and AKT1 enhances the phosphorylation and activation of AKT1, which, through an AKT1/GSK-3 β /NFATc1 signaling cascade, results in the increased production of the proapoptotic hormone Fas ligand and thus an increase in apoptosis.

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (rat ovary); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FTS) Polyclonal Antibody, Unconjugated (bs-13223R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.