

bs-22170R**[Primary Antibody]****BNC1 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 646	SWISS: Q01954	IF (1:100-500)
Target: BNC1		Reactivity: Mouse, Rat (predicted: Human)
Immunogen: KLH conjugated synthetic peptide derived from human BNC1: 211-310/994.		
Purification: affinity purified by Protein A		Predicted MW.: 111 kDa
Concentration: 1mg/ml		Subcellular Location: Cytoplasm ,Nucleus
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The zinc finger protein, basонуclin, is a putative rDNA transcription factor with highly restricted tissue distribution. Basонуclin is abundantly expressed in keratinocytes of the basal layer of the epidermis, the outer sheath of hair follicles and in the germ cells of the testis and ovary. Although basонуclin is associated with chromatin throughout the cell cycle, including mitosis, it disappears when cells become post-mitotic. In the epidermis, basонуclin, which is mainly localized to the cytoplasm, translocates to basal cell nuclei during different stages of keratinocyte growth. Basонуclin may enhance rRNA synthesis by elevating transcription from an rDNA promoter and inhibiting RNA polymerase I transcription through its zinc finger domain. Therefore, basонуclin may be a cell-type-specific transcription factor for rDNA transcription.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (BNC1) Polyclonal Antibody, Unconjugated (bs-22170R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (BNC1) Polyclonal Antibody, Unconjugated (bs-22170R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.