bs-19061R

- DATASHEET ----

[Primary Antibody]


NDOR1 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500) IF (1:100-500)
GenelD: 27158	SWISS: Q9UHB4	
Target: NDOR1		Reactivity: Mouse (predicted: Human, Rat, Pig, Cow, Dog)
Immunogen: KLH conjugated syn 51-150/597.	nthetic peptide derived from human N	
Purification: affinity purified by Protein A		Predicted MW.: 67 kDa
Concentration: 1mg/ml		MW.: OF NEW
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Location: Cytoplasm
contains both flavi dinucleotide (FAD) catalyzes the trans FMN cofactors to p	an NADPH-dependent diflavin reduct n mononucleotide (FMN) and flavin ac binding domains. The encoded prote fer of electrons from NADPH through otential redox partners. Alternative sp cranscript variants. [provided by RefSe	denine in FAD and Jlicing

- VALIDATION IMAGES -



2012]

Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NDOR1) Polyclonal Antibody, Unconjugated (bs-19061R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.