

bs-11012R

[Primary Antibody]

FAM98A Rabbit pAb

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

DATASHEET

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

GeneID: 25940

SWISS: Q8NCA5

Target: FAM98A

Immunogen: KLH conjugated synthetic peptide derived from human FAM98A: 251-350/519.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Encoding more than 700 genes, chromosome 15 is made up of approximately 106 million base pairs and is about 3% of the human genome. Angelman and Prader-Willi syndromes are associated with loss of function or deletion of genes in the 15q11-q13 region. In the case of Angelman syndrome, this loss is due to inactivity of the maternal 15q11-q13 encoded UBE3A gene in the brain by either chromosomal deletion or mutation. In cases of Prader-Willi syndrome, there is a partial or complete deletion of this region from the paternal copy of chromosome 15. Tay-Sachs disease is a lethal disorder associated with mutations of the HEXA gene, which is encoded by chromosome 15. Marfan syndrome is associated with chromosome 15 through the FBN1 gene. The FAM98 gene product has been provisionally designated FAM98 pending further characterization.

Applications: WB (1:500-2000)

IHC-P (1:100-500)

IHC-F (1:100-500)


IF (1:100-500)

Flow-Cyt (1µg/Test)


Reactivity: Human, Mouse
(predicted: Rat, Pig, Sheep, Cow, Chicken, Horse)

Predicted MW.: 55 kDa


VALIDATION IMAGES




25 µg total protein per lane of various lysates (see on figure) probed with FAM98A polyclonal antibody, unconjugated (bs-11012R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded (Human breast carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAM98A) Polyclonal Antibody, Unconjugated (bs-11012R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAM98A) Polyclonal Antibody, Unconjugated (bs-11012R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: hela(blue), the cells were fixed

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice.. Isotype Control Antibody: Rabbit IgG(orange) ; Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA ; Primary Antibody Dilution: 1µg in 100 µL 1X PBS containing 0.5% BSA(green).

— SELECTED CITATIONS —

- **[IF=16.7]** Shin-Ichiro Kanno. et al.Armadillo domain of ARID1A directly interacts with DNA-PKcs to couple chromatin remodeling with nonhomologous end joining (NHEJ) pathway..NUCLEIC ACIDS RESEARCH.2025 Feb 27;53(5):gkaf150. WB,IF ;Human. 40087883