

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

FPRL1 Rabbit pAb

Catalog Number: bs-23765R

Target Protein: FPRL1
Concentration: 1mg/1ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)
Reactivity: Human, Mouse

Predicted MW: 39 kDa Entrez Gene: 2358

Swiss Prot: P25090

Source: KLH conjugated synthetic peptide derived from human FPRL1: 1-100/350.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: FPRL1 is a low affinity receptor to N-formyl-methionyl peptides, which are powerful

neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of


neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-

calcium second messenger system. The activation of LXA4R could result in an anti-


inflammatory outcome counteracting the actions of proinflammatory signals such as LTB4

(leukotriene B4). FPRL1 has been reported in phagocytes, monocytes, neutrophils, differentiated myeloid cells from bone marrow, granulocyte HL-60 cells, and synovial fibroblasts. ESTs have been isolated from blood, leukocyte, lung, and placenta libraries.


VALIDATION IMAGES



Sample: SW480(Human) Cell Lysate at 30 ug LOVO(Human) Cell Lysate at 30 ug Primary: Anti-FPRL1 (bs-23765R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 39 kD Observed band size: 39 kD



Sample: Lane 1: U937 (Human) Cell Lysate at 30 ug Lane 2: HL60 (Human) Cell Lysate at 30 ug Lane 3: Hela (Human) Cell Lysate at 30 ug Lane 4: HepG2 (Human) Cell Lysate at 30 ug Lane 5: SW480 (Human) Cell Lysate at 30 ug Lane 6: Lovo (Human) Cell Lysate at 30 ug Primary: Anti-FPRL1 (bs-23765R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 39 kD Observed band size: 35 kD



Sample: Bone (Mouse) Lysate at 40 ug Primary: Anti- FPRL1 (bs-23765R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 39 kD Observed band size: 39 kD

PRODUCT SPECIFIC PUBLICATIONS

[IF=6.6] Ana Paula Girol. et al. Annexin A1 Mimetic Peptide and Piperlongumine: Anti-Inflammatory Profiles in Endotoxin-Induced Uveitis. Cells-Basel. 2021 Nov;10(11):3170 IHC; Rat. 34831393

[IF=5.895] Hui Wang. et al. Effect of Chlorogenic Acid via Upregulating Resolvin D1 Inhibiting the NF-kB Pathway on Chronic Restraint Stress-Induced Liver Inflammation. J AGR FOOD CHEM. 2022;70(34):10532–10542 WB; Rat. 35975781