## bs-8998R

## [ Primary Antibody ]

## **ORC1L/ORC1** Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal	-	<b>IHC-F</b> (1:100-500)
GenelD: 4998	<b>SWISS:</b> Q13415	<b>ICC/IF</b> (1:100-500)
Target: ORC1L/ORC1		ELISA (1:5000-10000)
Immunogen: KLH conjugated synthetic peptide derived from human ORC1L/ORC1: 701-800/861.		<b>Reactivity:</b> (predicted: Human, Mouse, Rat, Pig, Sheep, Cow, Dog, Horse)
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Predicted on the
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		MW.: <sup>97 kDa</sup> Subcellular Location: <sup>Nucleus</sup>
<b>Background:</b> The initiation of DNA replication is a multi-step process that depends on the formation of pre-replication complexes, which trigger initiation (1). Among the proteins required for establishing these complexes are the origin recognition complex (ORC) proteins (1). ORC proteins bind specifically to origins of replication where they serve as scaffold for the assembly of additional initiation factors (1). Human ORC subunits 1-6 are expressed in the nucleus of proliferating cells and tissues, such as the testis (2). ORC1 and ORC2 are both expressed at equivalent concentrations throughout the cell cycle; however, only ORC2 remains stably bound to chromatin (3,4). ORC4 and ORC6 are also expressed constantly throughout the cell cycle (5,6). ORC2, ORC3, ORC4 and ORC5 form a core complex upon which ORC6 and ORC1 assemble (7,8). The formation of this core complex suggests that ORC proteins play a crucial role in the G1-S transition in mammalian cells (8).		