

bs-13266R**[Primary Antibody]****Galectin 10 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 1178 Target: Galectin 10 Immunogen: KLH conjugated synthetic peptide derived from human Galectin 10: 51-142/142. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Charcot-Leyden crystals are endogenous hexagonal bipyramidal crystals present in human tissues and secretions. Presence of Charcot-Leyden crystals correlates with the increased numbers of peripheral blood or tissue eosinophils that occurs with parasitic and allergic processes. Galectin-10, also referred to as Charcot-Leyden crystal (CLC) protein, singularly makes up these crystals. Galectin-10, a member of the galectin family of beta-Galactoside binding proteins that bind to S-type animal lectins, is expressed solely in eosinophilic and basophilic leukocytes. Galectin-10 may possess carbohydrate or IgE-binding activities, and it plays a functional role in the biology of inflammation. Expression of galectin-10 is transcriptionally induced by butyric acid.	Isotype: IgG SWISS: Q05315	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
		Reactivity: Human, Rat
		Predicted MW.: 16 kDa
		Subcellular Location: Secreted ,Cytoplasm

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Galectin 10) Polyclonal Antibody, Unconjugated (bs-13266R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Galectin 10) Polyclonal Antibody, Unconjugated (bs-13266R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.