bs-18128R

[Primary Antibody]

phospho-IKB alpha (Ser32) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 4792 **SWISS:** P25963

Target: IKB alpha (Ser32)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

IKB alpha around the phosphorylation site of Ser32: D(p-S)GL.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes a member of the NF-kappa-B inhibitor family,

which contain multiple ankrin repeat domains. The encoded protein interacts with REL dimers to inhibit NF-kappa-B/REL complexes which are involved in inflammatory responses. The encoded protein moves between the cytoplasm and the nucleus via a nuclear localization signal and CRM1-mediated nuclear export. Mutations in this gene have been found in ectodermal dysplasia anhidrotic with T-cell immunodeficiency autosomal

dominant disease. [provided by RefSeq, Aug 2011]

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500)

Reactivity: Human, Mouse, Rat

(predicted: Rabbit, Pig,


Sheep, Cow)

Predicted ,


MW.: ^{36 kDa}

Subcellular Location: Cytoplasm ,Nucleus


VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-IKB alpha (Ser32)) Polyclonal Antibody, Unconjugated (bs-18128R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat lung); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-IKB alpha (Ser32)) Polyclonal Antibody, Unconjugated (bs-18128R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Tissue/cell:Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Antibody incubation with (phospho-IKB alpha (Ser32)) polyclonal Antibody, Unconjugated (bs-18128R) 1:100, 90 minutes at 37°C; followed by a FITC conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

- SELECTED CITATIONS -

- [IF=5.235] Pu, Luya. et al. Laminar shear stress alleviates monocyte adhesion and atherosclerosis development via miR-29b-3p/CX3CL1 axis regulation. J CELL SCI. 2022 Jul;135(14): WB; Human. 35735031
- [IF=5.195] Xiao Wang. et al. Kallistatin attenuates inflammatory response in rheumatoid arthritis via the NF-κB signaling pathway. EUR J PHARMACOL. 2023 Jan;:175530 WB; Mouse. 36690053
- [IF=3.776] Yang Yang. et al. The NF-κB pathway plays a vital role in rat salivary gland atrophy model. HELIYON. 2023 Mar; 9(3): e14288 WB; Rat. 36950625

- [IF=3.845] Peng LY et al. Protective Effect of Piceatannol Against Acute Lung Injury Through Protecting the Integrity of Air-Blood Barrier and Modulating the TLR4/NF-kB Signaling Pathway Activation. Front Pharmacol. 2020 Jan 22;10:1613. WB; Mouse. 32038265
- [IF=3.1] Meng Yan. et al. Diammonium Glycyrrhizinate Inhibited Inflammatory Response and Modulated Serum Metabolism in Poly(I:C)-induced Pneumonia Model Mice. SHOCK. 2024 Mar;:10.1097/SHK.00000000000002353 WB; Mouse. 38526139