bs-13278R

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

GAMT Rabbit pAb

- DATASHEET -

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

GeneID: 2593 **SWISS:** Q14353

Target: GAMT

Immunogen: KLH conjugated synthetic peptide derived from human GAMT:

101-200/236.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: In the creatine biosynthesis pathway, glycine is converted to guanidinoacetate by amidinotransferase, and guanidinoacetate is

then converted to creatine by Guanidinoacetate N-methyltransferase (GAMT). GAMT, a methyltransferase, uses S-adenosylmethionine as the methyl donor for this reaction.

Methyltransferases are a type of transferase enzyme which transfers a methyl group to nucleic bases in DNA or amino acids in protein. Encoding a 236 amino acid protein, the human GAMT gene maps to chromosome 19p13.3. Defects in the GAMT gene leads to GAMT deficiency, which is associated with guanidinoacetate accumulation and decreased levels of creatine excretion in brain.

Such biochemical changes are thought to lead to various neurological syndromes and muscular hypotonia.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500)


Reactivity: Human, Mouse, Rat

(predicted: Pig, Dog)


Predicted MW.: ^{26 kDa}

Subcellular Location: Cytoplasm ,Nucleus


VALIDATION IMAGES



Tissue/cell: Human hepatocellular carcinoma; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-GMAT Polyclonal Antibody, Unconjugated(bs-13278R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (GAMT) Polyclonal Antibody, Unconjugated (bs-13278R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



HepG2 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (GAMT) polyclonal Antibody, Unconjugated (bs-13278R) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.