### bs-13280R

## [ Primary Antibody ]

# RASA1 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)

Reactivity: Rat (predicted: Human, Mouse, Pig, Sheep, Cow, Chicken, Dog, Horse)

Predicted MW.: 116 kDa

Subcellular Location: Cytoplasm

Clonality: Polyclonal

Host: Rabbit

- DATASHEET -

SWISS: P20936

Isotype: IgG

GenelD: 5921 Target: RASA1

Immunogen: KLH conjugated synthetic peptide derived from human Ras GTPase-activating protein 1: 451-550/1047.


Purification: affinity purified by Protein A

### Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** The mammalian c-H-, c-K- and N-Ras proto-oncogenes encode ubiquitously expressed proteins (1,2). p21Ras can exist in either a physiologically quiescent GDP-binding state or a GTP-binding signal-emitting state (3). Oncogenic p21Ras proteins are trapped in the excited signal-emitting state because the mechanism normally employed to delimit their excitation period, hydrolysis of their bound GTP to GDP, is impaired as a result of specific mutations (3). Interaction of p21Ras with GTPase activating protein (GAP) can increase hydrolysis of p21Ras-bound GTP by as much as 1000-fold (4,5). The product of the neurofibromatosis type 1 gene (NF1) has also been shown to exhibit p21Ras GAP activity (6,7), and proteins that stimulate the GTPase activity of three other low molecular weight GTPases, including Rho, Rab 3A and Rap 1, have also been described (8,9).

#### – VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RASA1) Polyclonal Antibody, Unconjugated (bs-13280R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.